

# On the Kolmogorov-Smirnov Test for Normality with Mean and Variance

Unknown

Hubert W. Lilliefors

*Journal of the American Statistical Association*, Vol. 62, No. 318.  
(Jun., 1967), pp. 399-402.

Stable URL:

<http://links.jstor.org/sici?sici=0162-1459%28196706%2962%3A318%3C399%3AOTKTFN%3E2.0.CO%3B2-G>

*Journal of the American Statistical Association* is currently published by American Statistical Association.

TABLE 1. TABLE OF CRITICAL VALUES OF  $D$

The values of  $D$  given in the table are critical values associated with selected values of  $N$ . Any value of  $D$  which is greater than or equal to the tabulated value is significant at the indicated level of significance. These values were obtained as a result of Monte Carlo calculations, using 1,000 or more samples for each value of  $N$ .

Sample Size $N$	Level of Significance for $D = \text{Max}  F^*(X) - S_N(X) $				
	.20	.15	.10	.05	.01
4	.300	.319	.352	.381	.417
5	.285	.299	.315	.337	.405
6	.265	.277	.294	.319	.364
7	.247	.258	.276	.300	.348
8	.233	.244	.261	.285	.331
9	.223	.233	.249	.271	.311
10	.215	.224	.239	.258	.294
11	.206	.217	.230	.249	.284
12	.199	.212	.223	.242	.275
13	.190	.202	.214	.234	.268
14	.183	.194	.207	.227	.261
15	.177	.187	.201	.220	.257
16	.173	.182	.195	.213	.250
17	.169	.177	.189	.206	.245
18	.166	.173	.184	.200	.239
19	.163	.169	.179	.195	.235
20	.160	.166	.174	.190	.231
25	.149	.153	.165	.180	.203
30	.131	.136	.144	.161	.187
Over 30	$\frac{.736}{\sqrt{N}}$	$\frac{.768}{\sqrt{N}}$	$\frac{.805}{\sqrt{N}}$	$\frac{.886}{\sqrt{N}}$	$\frac{1.031}{\sqrt{N}}$